

Dríva™ Drills Gold Series

Usage Chart

For use with TSX°, Trabecular Metal°, TSV°, and Eztetic° Implants



Use a dry erase marker to track the total number of osteotomy preparations that are performed with each drill. It is recommended that reusable drills be replaced after 15 uses.* Single-use drills must be disposed of after each procedure.

			7	7		7		7	7	7	ħ	7			7	Ž	7				7		7	7	7
	0201G 2.1/1.6 mmD,	TSV23G 2.3 mmD	EZT24G 2.4 mmD	EZT28D24G 2.8/2.4 mmD	TSV28G 2.8 mmD	TSV34D28G 3.4/2.8 mmD	TSV34G 3.8 mmD	TSV38D34G 3.8/3.4 mmD	TSV38G 3.8 mmD	TSV44D38G 4.4/3.8 mmD	TSV44G 4.4 mmD	TSV51D44G 5.1/4.4 mmD	TSV51G 5.1 mmD	TSV57D51G 5.7/5.1 mmD	TSV23SG	TSV28SG	TSV34D28SG 3.4/2.8 mmD	TSV34SG	TSV38D34SG 3.8/3.4 mmD	TSV38SG	TSV44D38SG 4.4/3.8 mmD	TSV44SG	TSV51D44SG 5.1/4.4 mmD	TSV51SG	TSV57D51SG 5.7/5.1 mmD
	Tapered Pilot Drill 8.0 mm - 11.5 mmL	Drill, 22 mmL	Drill, 22 mmL	Step Drill, 22 mmL	Drill, 22 mmL	Step Drill, 22 mmL	Drill, 22 mmL	Step Drill, 22 mmL	Drill, 22 mmL	Step Drill, 22 mmL	Drill, 22 mmL	Step Drill, 22 mmL	Drill, 22 mmL	Step Drill, 22 mmL	2.3 mmD Drill, 16 mmL	2.8 mmD Drill, 16 mmL	Step Drill, 16 mmL	3.4 mmD Drill, 16 mmL	Step Drill, 16 mmL	3.8 mmD Drill, 16 mmL	Step Drill, 16 mmL	4.4 mmD Drill, 16 mmL	Stop Drill 16	5.1 mmD Drill, 16 mmL	Step Drill, 16 mmL
Drilling Sequence																									
Time to Order New Drills	1	1	1	1	1	1	1	0	1	1	1	0	1	1	1	1	1	1	0	1	1	1	0	1	1
	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6
	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
	11)	11)	11)	11	11)	11)	11)	•	11)	11)	11)	0	11)	11	11)	11)	11)	11)	•	11)	11)	11)	•	11)	11)
	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12
	13)	13)	13	13	13	13	13	13	13	13	13	13	13	13	13)	13	13	13	13	13	13	13	13	13)	13
	14)	14)	14)	14)	14)	14)	14)	14	14)	14)	14)	14	14)	14)	14)	14)	14)	14)	14	14)	14)	14)	14	14)	14)
	15)	15)	15)	15	15)	15)	15)	15	15)	15)	15)	15	15)	15	15)	15)	15)	15)	15	15)	15)	15)	15	15)	15)

^{*} Single-use drills must be disposed of after each procedure. This chart is intended to be used as a reference guide only. The number of uses per drill will vary and depends on a variety of factors including bone density encountered, proper handling, and cleaning. Refer to Instructions for Use (IFU) for ZimVie Kits and Instruments (P-ZBDINSTRP) available at labeling.zimvie.com for complete cleaning and sterilization information.

NOTE: Surgical instruments are susceptible to damage for a variety of reasons including but not limited to: prolonged use, misuse, rough or improper handling, inadequate cleaning, etc. Care must be taken to avoid compromising the intended performance of the instrument.

For more information, visit ZimVie.com

ZimVie 4555 Riverside Drive Palm Beach Gardens, FL 33410 1-800-342-5454 Phone: +1-561-776-6700 Fax: +1-561-776-1272 Unless otherwise indicated, as referenced herein, all trademarks and intellectual property rights are the property of ZimVie Inc. or an affiliate; and all products are manufactured by one or more of the dental subsidiaries of ZimVie Inc. (Biomet 3i, LLC, Zimmer Dental, Inc., etc.) and marketed and distributed by ZimVie Dental and its authorized marketing partners. For additional product information, please refer to the individual product

labeling or instructions for use. Product clearance and availability may be limited to certain countries/regions. This material is intended for clinicians only and does not comprise medical advice or recommendations. Distribution to any other recipient is prohibited. This material may not be copied or reprinted without the express written consent of ZimVie. ZV1629 REV B 01/24 ©2024 ZimVie, Inc. All rights reserved.

